arXiv:1409.2074v1 [math.CA] 7 Sep 2014

Bilateral zeta functions associated with the multiple sine functions

Genki Shibukawa

Abstract

We introduce two types bilateral zeta functions, which are related to the primitive and normalized multiple sine functions respectively. Further, we establish their main properties, that is, Fourier expansions, analytic continuations, differential and difference equations, special values. By applying these results, we obtain not only some generalization of the primitive and normalized multiple sine functions but also simple construction of the multiple sine function theory.

1 Introduction

The primitive multiple sine functions $\mathscr{S}_r(z)$ $(r \in \mathbb{Z}_{\geq 1})$

$$\mathscr{S}_r(z) := \exp\left(\frac{z^{r-1}}{r-1}\right) \prod_{m=1}^{\infty} \left(P_r\left(\frac{z}{m}\right) P_r\left(-\frac{z}{m}\right)^{(-1)^{r-1}} \right)^{m^{r-1}} = \exp\left(\int_0^z \pi t^{r-1} \cot \pi t \, dt \right)$$

were introduced and studied by Hölder $(r = 2)$ and Kurokawa $(r \geq 3)$ (see [KK]). Here, we put

$$P_r(u) := (1-u) \exp\left(\sum_{j=1}^{r} \frac{u^j}{j} \right).$$

Moreover, Kurokawa has constructed a normalized multiple sine functions $S_r(z)$. This function is defined as

$$S_r(z) := \exp\left(-\frac{\partial \zeta_r}{\partial s}(0,z) + (-1)^r \frac{\partial \zeta_r}{\partial s}(0, r-z) \right).$$

Here, $\zeta_r(s,z)$ is the multiple Hurwitz function, which extends holomorphic function on the whole s-plane except for possible simple poles at $s = 1, \ldots, r$ under $\operatorname{Re} z > 0$.

$$\zeta_r(s,z) := \sum_{m_1,\ldots,m_r \geq 0} \frac{1}{(m_1 + \cdots + m_r + z)^s} = \frac{1}{(r-1)!} \sum_{m \geq 0} \frac{(m+1)_{r-1}}{(m+z)^s} \quad (\operatorname{Re} z > 0).$$

The intimate relation between these two kinds of multiple sine functions is established by Kurokawa (see [KK]).

Furthermore, for the normalized multiple sine functions $S_r(z)$, more generalization have been studied in Kurokawa-Wakayama [KW]. Actually, for any $n \in \mathbb{Z}_{\geq 1}$, a generalization of the normalized multiple sine functions $S_{r,n}(z)$ is given by

$$S_{r,n}(z) := \exp\left(-\frac{\partial \zeta_r}{\partial s}(1-n,z) + (-1)^{r+n-1}\frac{\partial \zeta_r}{\partial s}(1-n,r-z)\right).$$

We remark $S_{r,1}(z) = S_r(z)$. On the other hand, for the primitive type, such a kind of the generalization has not been studied yet.

In this article, we introduce two types bilateral zeta functions

$$H_r(s,z) := \sum_{m\in\mathbb{Z}} \frac{m^{r-1}}{(m+z)^s}$$

and

$$K_r(s,z) := \frac{1}{(r-1)!}\sum_{m\in\mathbb{Z}} \frac{(m+1)_{r-1}}{(m+z)^s}.$$

Since the derivatives at $s=0$ of $H_r(s,z)$ and $K_r(s,z)$ are equal to the primitive and normalized multiple sine functions up to some exponential factors, we define the generalized multiple sine functions by a derivation of $H_r(s,z)$ and $K_r(s,z)$ at $s = 0, -1, -2, \ldots$. From this point of view, we are not only succeed in providing a generalization of the primitive multiple sine function, but also in providing simple construction of the multiple sine function theory.

In Section 2, we give their fundamental properties, that is, Fourier expansions, analytic continuations, differential and difference equations, special values. In Section 3, by applying results of Section 2, we introduce a generalization of the primitive and normalized multiple sine functions and establish their basic properties.

2 Definition and basic properties

Throughout the paper, we denote the ring of rational integers by $\mathbb{Z}$, the field of real numbers by $\mathbb{R}$, the field of complex numbers by $\mathbb{C}$ and $i := \sqrt{-1}$. Further, we fix the branch of any complex numbers c.

$$-\pi < \arg c \leq \pi.$$

In particular, we define $\arg 0 := 0$. We assume $r \in \mathbb{Z}_{\geq 2}, n \in \mathbb{Z}_{\geq 1}, s \in \mathbb{C}$ and $z \in \mathbb{C}$ with $\operatorname{Im} z \geq 0$ unless otherwise specified.

Definition 2.1. Let $\operatorname{Re} s > r$.
(1) We define a bilateral zeta function associated with the primitive multiple sine functions by the series

$$H_r(s,z) := \sum_{m\in\mathbb{Z}} \frac{m^{r-1}}{(m+z)^s}. \tag{2.1}$$

(2) We also introduce a bilateral zeta function associated with the normalized multiple sine functions as

$$K_r(s,z) := \frac{1}{(r-1)!}\sum_{m\in\mathbb{Z}}\frac{(m+1)_{r-1}}{(m+z)^s}. \tag{2.2}$$

Here, $(X)_r$ is the shifted factorial defined by

$$(X)_r := \begin{cases} 1 & (r=0) \\ X(X+1)\cdot(X+r-1) & (\text{otherwise}) \end{cases}.$$

In particular, we put

$$\xi(s,z) := H_1(s,z) = K_1(s,z). \tag{2.3}$$

These series absolutely converge for $\operatorname{Re} s > r$. Moreover, from the following well-known formula, $\xi(s,z)$ is continued analytically to $\mathbb{C}$ as a holomorphic function in s.

Lemma 2.2. (1) *If* $1 > \operatorname{Re} z > 0, \operatorname{Im} z \geq 0$, *then*

$$\xi(s,z) = \zeta_1(s,z) + e^{-\pi i s}\zeta_1(s,1-z). \tag{2.4}$$

By analytic continuation of the Hurwitz zeta function $\zeta_1(s,z)$, $\xi(s,z)$ *extends analytically as a holomorphic function on the whole* s*-plane under the conditions* $1 > \operatorname{Re} z > 0, \operatorname{Im} z \geq 0$.
(2) *(See [AAR] Chapter II. Exercise 37) If* $\operatorname{Im} z > 0$, *then*

$$\xi(s,z) = \frac{(2\pi)^s}{\Gamma(s)}e^{-\frac{\pi}{2}is}\operatorname{Li}_{1-s}(e^{2\pi i z}). \tag{2.5}$$

Here, $\operatorname{Li}_\alpha(X)$ *is the polylogarithm*

$$\operatorname{Li}_\alpha(X) := \sum_{m=1}^{\infty}\frac{X^m}{m^\alpha}.$$

From this expression, $\xi(s,z)$ *extends analytically as a holomorphic function on the whole* s*-plane under the conditions* $\operatorname{Im} z > 0$.

By this fundamental formula, we also more results for $\xi(s,z)$, which are needed in later.

Corollary 2.3. (1) *For any* $s \in \mathbb{C}$, *we have*

$$\xi(s,z+1) = \xi(s,z). \tag{2.6}$$

(2)

$$\xi(1,z) = \pi i + \pi\cot\pi z, \tag{2.7}$$

$$\xi(n+1,z) = \frac{\pi^{n+1}}{n!}\sum_{k=0}^{n-1}\left\langle{n \atop k}\right\rangle(\cot\pi z - i)^{k+1}(\cot\pi z + i)^{n-k} \tag{2.8}$$

$$= \frac{1}{n!}\sum_{k=0}^{n-1}\sum_{l=0}^{k+1}\left\langle{n \atop k}\right\rangle\binom{k+1}{l}(-2\pi i)^l\xi(1,z)^{n+1-l}, \tag{2.9}$$

$$\xi(1-n,z) = 0. \tag{2.10}$$

Here, $\left\langle {n \atop k} \right\rangle$ *are Eulerian numbers which are the number of permutations of the set* $\{1, \ldots, n\}$ *having* k *permutation ascents.*
(3) *For any* $n \in \mathbb{Z}_{\geq 1}$,

$$\frac{\partial \xi}{\partial s}(1-n, z) = \frac{(n-1)!}{(2\pi i)^{n-1}} \mathrm{Li}_n(e^{2\pi i z}). \tag{2.11}$$

Proof. With the exception of (2.8) and (2.9), all properties are prove immediately from Lemma 2.2. (2.8) follows from the theorem of G. Rzadkowski [R] that is if there exist some constants a, α, β such that

$$\frac{d}{dz} y(z) = a(y(z) - \alpha)(y(z) - \beta),$$

then

$$y^{(n)}(z) = a^n \sum_{k=0}^{n-1} \left\langle {n \atop k} \right\rangle (y(z) - \alpha)^{k+1} (y(z) - \beta)^{n-k}.$$

Actually, we remark

$$\frac{d}{dz} \xi(1, z) = \frac{d}{dz} \pi \cot \pi z = -\pi^2 (\cot \pi z - i)(\cot \pi z + i)$$

and

$$\frac{d}{dz} \xi(n, z) = -n \xi(n+1, z).$$

Therefore,

$$\xi(n+1, z) = \frac{(-1)^n}{n!} \frac{d^n}{d^n z} \pi \cot \pi z = \frac{\pi^{n+1}}{n!} \sum_{k=0}^{n-1} \left\langle {n \atop k} \right\rangle (\cot \pi z - i)^{k+1} (\cot \pi z + i)^{n-k}.$$

(2.9) follows from (2.7) and (2.8). □

The relation between $H_r(s, z)$ and $K_r(s, z)$ is given by the following Proposition.

Proposition 2.4. (1) *Let* $\left\{ {r \atop k} \right\}$ *be the Stirling numbers of the second kind. We have*

$$H_r(s, z) = \sum_{k=1}^{r} (-1)^{r-k} (k-1)! \left\{ {r \atop k} \right\} K_k(s, z). \tag{2.12}$$

(2) *Let* $\left[{r \atop k} \right]$ *be the Stirling numbers of the first kind. We have*

$$K_r(s, z) = \frac{1}{(r-1)!} \sum_{k=1}^{r} \left[{r \atop k} \right] H_k(s, z). \tag{2.13}$$

Proof. We recall the properties of the Stirling numbers

$$m^r = \sum_{k=1}^{r} (-1)^{r-k} \left\{ {r \atop k} \right\} (m)_k = m \sum_{k=1}^{r} (-1)^{r-k} \left\{ {r \atop k} \right\} (m+1)_{k-1},$$
$$(m)_r = m(m+1)_{r-1} = \sum_{k=1}^{r} \left[{r \atop k} \right] m^k = m \sum_{k=1}^{r} \left[{r \atop k} \right] m^{k-1}.$$

From these properties, we have

$$H_r(s,z) = \sum_{m\in\mathbb{Z}} \frac{1}{(m+z)^s} \sum_{k=1}^{r} (-1)^{r-k} \left\{ {r \atop k} \right\} (m+1)_{k-1} = \sum_{k=1}^{r} (-1)^{r-k} (k-1)! \left\{ {r \atop k} \right\} K_k(s,z),$$
$$K_r(s,z) = \frac{1}{(r-1)!} \sum_{m\in\mathbb{Z}} \frac{1}{(m+z)^s} \sum_{k=1}^{r} \left[{r \atop k} \right] m^{k-1} = \frac{1}{(r-1)!} \sum_{k=1}^{r} \left[{r \atop k} \right] H_k(s,z).$$

□

Proposition 2.5 ($\xi(s,z)$ expressions)**.** (1)

$$H_r(s,z) = \sum_{k=0}^{r-1} \binom{r-1}{k} (-z)^{r-1-k} \xi(s-k,z). \tag{2.14}$$

(2)

$$K_r(s,z) = \frac{1}{(r-1)!} \sum_{k=1}^{r} \left[{r \atop k} \right] \sum_{l=0}^{k-1} \binom{k-1}{l} (-z)^{k-l-1} \xi(s-l,z). \tag{2.15}$$

Proof. (1) By the binomial theorem,

$$\begin{aligned} H_r(s,z) &= \sum_{m\in\mathbb{Z}} \frac{(m+z-z)^{r-1}}{(m+z)^s} \\ &= \sum_{m\in\mathbb{Z}} \frac{1}{(m+z)^s} \sum_{k=0}^{r-1} \binom{r-1}{k} (-z)^{r-1-k} (m+z)^k \\ &= \sum_{k=0}^{r-1} \binom{r-1}{k} (-z)^{r-1-k} \xi(s-k,z). \end{aligned}$$

(2) This result follows from (2.13) and (2.14) immediately. □

From Lemma 2.2 and Proposition 2.5, the functions $H_r(s,z)$ and $K_r(s,z)$ are analytically continued to $s \in \mathbb{C}$ as holomorphic functions. Therefore, although all results in this section hold for all s, we only need to prove them for large enough $\operatorname{Re} s$.

Further, from Corollary 2.3 and Proposition 2.5, we obtain some special values of $H_r(s,z)$ and $K_r(s,z)$.

Corollary 2.6 (special values)**.** *Let $n \geq 1$ and $r-1 \geq p \geq 1$.*
(1)

$$H_r(1,z) = (-z)^{r-1}(\pi i + \pi \cot \pi z), \tag{2.16}$$

$$\begin{aligned} H_r(1+p,z) = {} & \binom{r-1}{p}(-z)^{r-p-1}(\pi i + \pi \cot \pi z) + \sum_{k=0}^{p-1} \binom{r-1}{k}(-z)^{r-1-k}\frac{\pi^{p+1-k}}{(p-k)!} \\ & \cdot \sum_{l=0}^{p-k-1} \left\langle {p-k \atop l} \right\rangle (\cot \pi z - i)^{l+1}(\cot \pi z + i)^{p-k-l}, \end{aligned} \tag{2.17}$$

$$\begin{aligned} H_r(r+n,z) = {} & \sum_{k=0}^{r-1} \binom{r-1}{k}(-z)^{r-1-k}\frac{\pi^{r+n-k}}{(r+n-k-1)!} \\ & \cdot \sum_{l=0}^{r+n-k-2} \left\langle {r+n-k-1 \atop l} \right\rangle (\cot \pi z - i)^{l+1}(\cot \pi z + i)^{r+n-k-1-l}, \end{aligned} \tag{2.18}$$

$$H_r(1-n,z) = 0. \tag{2.19}$$

(2)

$$K_r(1,z) = \frac{(1-z)_{r-1}}{(r-1)!}(\pi i + \pi \cot \pi z), \tag{2.20}$$

$$\begin{aligned} K_r(1+p,z) = {} & \sum_{k=1}^{p} \begin{bmatrix} r \\ k \end{bmatrix} \sum_{l=0}^{k-1} \binom{k-1}{l}(-z)^{k-1-l}\frac{\pi^{1+p-l}}{(p-l)!} \\ & \cdot \sum_{q=0}^{p-l-1} \left\langle {p-l \atop q} \right\rangle (\cot \pi z - i)^{q+1}(\cot \pi z + i)^{p-l-q} \\ & + \frac{1}{(r-1)!} \sum_{k=p+1}^{r} \begin{bmatrix} r \\ k \end{bmatrix} \sum_{l=0}^{k-1} \binom{k-1}{p}(-z)^{k-p-1}(\pi i + \pi \cot \pi z) \\ & + \frac{1}{(r-1)!} \sum_{k=p+1}^{r} \begin{bmatrix} r \\ k \end{bmatrix} \sum_{l=0}^{p-1} \binom{k-1}{l}(-z)^{k-l-1}\frac{\pi^{p+1-l}}{(p-l)!} \\ & \cdot \sum_{q=0}^{p-l-1} \left\langle {p-l \atop q} \right\rangle (\cot \pi z - i)^{q+1}(\cot \pi z + i)^{p-l-q}, \end{aligned} \tag{2.21}$$

$$\begin{aligned} K_r(r+n,z) = {} & \frac{1}{(r-1)!} \sum_{k=1}^{r} \begin{bmatrix} r \\ k \end{bmatrix} \sum_{l=0}^{k-1} \binom{k-1}{l}(-z)^{k-1-l}\frac{\pi^{r+n-l}}{(r+n-l-1)!} \\ & \cdot \sum_{q=0}^{r+n-l-2} \left\langle {r+n-l-1 \atop q} \right\rangle (\cot \pi z - i)^{q+1}(\cot \pi z + i)^{r+n-l-1-q}, \end{aligned} \tag{2.22}$$

$$K_r(1-n,z) = 0. \tag{2.23}$$

We also have some expression of $H_r(s,z)$ and $K_r(s,z)$ by using the multiple Hurwitz zeta functions.

Proposition 2.7 ($\zeta_r(s,z)$ expressions)**.** (1) *If* $1 > \mathrm{Re}\, z > 0$, $\mathrm{Im}\, z \geq 0$, *then for any* $s \in \mathbb{C}$, *we have*

$$H_r(s,z) = \sum_{k=1}^{r} (-1)^{r-k}(k-1)! \left\{ {r \atop k} \right\} (\zeta_k(s,z) + (-1)^{k-1} e^{-\pi i s} \zeta_k(s, k-z)). \tag{2.24}$$

(2) *If* $r > \mathrm{Re}\, z > 0$, $\mathrm{Im}\, z \geq 0$, *then for any* $s \in \mathbb{C}$, *we have*

$$K_r(s,z) = \zeta_r(s,z) + (-1)^{r-1} e^{-\pi i s} \zeta_r(s, r-z) \tag{2.25}$$

$$= z^{-s} + \zeta_r(s, 1+z) + (-1)^{r-1} e^{-\pi i s} \zeta_r(s, r-z). \tag{2.26}$$

(3) *If* $1 > \mathrm{Re}\, z > -1$, $\mathrm{Im}\, z \geq 0$, *then for any* $s \in \mathbb{C}$, *we have*

$$H_r(s,z) = \sum_{k=0}^{r-1} \binom{r-1}{k} (-z)^{r-1-k} (\zeta(s-k, 1+z) + (-1)^{k-1} e^{-\pi i s} \zeta(s-k, 1-z)). \tag{2.27}$$

Proof. (1) (2.24) follows from (2.25) and (2.12).
(2) We decompose the sum $K_r(s,z)$ as follows.

$$K_r(s,z) = \frac{1}{(r-1)!} \sum_{m \geq 0} \frac{(m+1)_{r-1}}{(m+z)^s} + \frac{1}{(r-1)!} \sum_{m \geq 0} \frac{(-m)_{r-1}}{(-m-1+z)^s}.$$

Here, we remark $0 < \arg(-m-1+z) < \pi$ and

$$\sum_{m \geq 0} \frac{(-m)_{r-1}}{(-m-1+z)^s} = \sum_{m \geq r-1} \frac{(-m)_{r-1}}{(-m-1+z)^s} = (-1)^{r-1} e^{-\pi i s} \sum_{m \geq 0} \frac{(m+1)_{r-1}}{(m+r-z)^s}.$$

From this calculation, we have (2.25). (2.26) follows from (2.25) and the definition of $\zeta_r(s,z)$.
(3) Since $r \geq 2$ and $0 < \arg(-m-1+z) < \pi$ under the condition,

$$\begin{aligned}
H_r(s,z) &= \sum_{m \neq 0} \frac{m^{r-1}}{(m+z)^s} \\
&= \sum_{m \geq 0} \frac{(m+1+z-z)^{r-1}}{(m+1+z)^s} + (-1)^{r-1} e^{-\pi i s} \sum_{m \geq 0} \frac{(m+1-z+z)^{r-1}}{(m+1-z)^s} \\
&= \sum_{k=0}^{r-1} \binom{r-1}{k} (-z)^{r-1-k} \zeta(s-k, 1+z) \\
&\quad + (-1)^{r-1} e^{-\pi i s} \sum_{k=0}^{r-1} \binom{r-1}{k} z^{r-1-k} \zeta(s-k, 1-z) \\
&= \sum_{k=0}^{r-1} \binom{r-1}{k} (-z)^{r-1-k} (\zeta(s-k, 1+z) + (-1)^{k-1} e^{-\pi i s} \zeta(s-k, 1-z)).
\end{aligned}$$

□

In addition, by applying the Fourier expansion for $\xi(s,z)$, we obtain the following Fourier expansions of H and K.

Proposition 2.8 (Fourier expansions)**.** *Let* $\operatorname{Im} z > 0$.
(1) *For any* $s \in \mathbb{C}$,

$$H_r(s,z) = \frac{(2\pi)^s}{\Gamma(s)} e^{-\frac{\pi}{2}is}(-z)^{r-1} \sum_{k=0}^{r-1} \frac{(1-r)_k (1-s)_k}{k!} (2\pi i z)^{-k} \mathrm{Li}_{1+k-s}(e^{2\pi i z}). \tag{2.28}$$

(2)

$$K_r(s,z) = \frac{(2\pi)^s}{\Gamma(s)} \frac{e^{-\frac{\pi}{2}is}}{(r-1)!} \sum_{k=1}^{r} \begin{bmatrix} r \\ k \end{bmatrix} (-z)^{k-1} \sum_{l=0}^{k-1} \frac{(1-k)_l (1-s)_l}{l!} (2\pi i z)^{-l} \mathrm{Li}_{1+l-s}(e^{2\pi i z}). \tag{2.29}$$

Applying Proposition 2.8, we obtain special values of derivation of H and K.

Proposition 2.9. (1) *If* $\operatorname{Im} z > 0$ *or* $1 > z > -1$*, then for any* $n \in \mathbb{Z}_{\geq 1}$ *and* $s \in \mathbb{C}$,

$$\frac{\partial H_r}{\partial s}(1-n,z) = \sum_{k=0}^{r-1} \frac{(n+k-1)!}{(2\pi i)^{n+k-1}} \binom{r-1}{k} (-z)^{r-1-k} \mathrm{Li}_{n+k}(e^{2\pi i z}). \tag{2.30}$$

(2) *If* $\operatorname{Im} z > 0$ *or* $r > z > 0$*, for any* $n \in \mathbb{Z}_{\geq 1}$ *and* $s \in \mathbb{C}$,

$$\frac{\partial K_r}{\partial s}(1-n,z) = \frac{1}{(r-1)!} \sum_{k=1}^{r} \begin{bmatrix} r \\ k \end{bmatrix} \sum_{l=0}^{k-1} \binom{k-1}{l} \frac{(n+l-1)!}{(2\pi i)^{n+l-1}} (-z)^{k-l-1} \mathrm{Li}_{n+l}(e^{2\pi i z}). \tag{2.31}$$

Moreover, if $n > 1$*, this result holds for* $\operatorname{Im} z > 0$ *or* $r > z \geq 0$.

Proof. If $\operatorname{Im} z > 0$, then (2.30) and (2.31) follow from Proposition 2.8 immediately. Further, we remark the right hand sides of (2.30) and (2.31) is well-defined under the conditions. Thus, it is enough to show the well-definedness of the left hand sides of (2.30) and (2.31).
(1) By (2.27), $H_r(s,z)$ is holomorphic function of s under $\operatorname{Im} z \geq 0$, $1 > \operatorname{Re} z > -1$. Hence, by identity theorem, we have the conclusion.
(2) By (2.25), $K_r(s,z)$ is an entire function for s under $\operatorname{Im} z \geq 0$, $r > \operatorname{Re} z > 0$. Thus, by identity theorem, we have (2.31) under $\operatorname{Im} z > 0$ or $r > z > 0$.

On the other hands, by (2.26), the left hand sides of (2.31) has the following expression.

$$\begin{aligned} \frac{\partial H_r}{\partial s}(1-n,z) = &-z^{n-1} \log z - \pi i (-1)^{r+n} \zeta_r(1-n, r-z) \\ &+ \frac{\partial \zeta_r}{\partial s}(1-n, 1+z) + \frac{\partial \zeta_r}{\partial s}(1-n, r-z). \end{aligned}$$

When $n > 1$, from the above expression of $\frac{\partial H_r}{\partial s}(1-n,z)$ is entire in $\operatorname{Im} z \geq 0$, $r > \operatorname{Re} z \geq 0$. Therefore, by identity theorem, we obtain the conclusion.

□

Corollary 2.10. (1)

$$\frac{\partial H_r}{\partial s}(1-n,0) = \frac{(n+r-2)!}{(2\pi i)^{n+r-2}}\zeta(n+r-1), \tag{2.32}$$

$$\frac{\partial K_r}{\partial s}(-n,0) = \frac{1}{(r-1)!}\sum_{k=1}^{r}\begin{bmatrix} r \\ k \end{bmatrix}\frac{(n+k-1)!}{(2\pi i)^{n+k-1}}\zeta(n+k). \tag{2.33}$$

Here, $\zeta(s) := \zeta_1(s,1)$ *is the Riemann zeta function.*
(2)

$$\frac{\partial H_r}{\partial s}\left(1-n,\frac{1}{2}\right) = \sum_{k=0}^{r-1}\frac{(n+k-1)!}{(2\pi i)^{n+k-1}}\binom{r-1}{k}(-2)^{-r+1+k}(2^{1-n-k}-1)\zeta(n+k), \tag{2.34}$$

$$\frac{\partial K_r}{\partial s}\left(1-n,\frac{1}{2}\right) = \frac{1}{(r-1)!}\sum_{k=1}^{r}\begin{bmatrix} r \\ k \end{bmatrix}\sum_{l=0}^{k-1}\binom{k-1}{l}\frac{(n+l-1)!}{(2\pi i)^{n+l-1}}(-2)^{-k+l+1}(2^{1-n-l}-1)\zeta(n+l). \tag{2.35}$$

We remark $\lim_{n+k\to 1}(2^{1-n-k}-1)\zeta(n+k) = -\log 2$.

Proposition 2.11 (difference equations for z)**.** *Let* $l \in \mathbb{Z}_{\geq 1}$.
(1)

$$H_r(s,z+l) = \sum_{k=0}^{r-1}(-l)^k\binom{r-1}{k}H_{r-k}(s,z). \tag{2.36}$$

(2)

$$K_r(s,z+l) = K_r(s,z) - \sum_{k=0}^{l-1}K_{r-1}(s,z+k). \tag{2.37}$$

Proof. (1)

$$H_r(s,z+l) = \sum_{m\in\mathbb{Z}}\frac{m^{r-1}}{(m+z+l)^s} = \sum_{m\in\mathbb{Z}}\frac{(m-l)^{r-1}}{(m+z)^s} = \sum_{m\in\mathbb{Z}}\frac{1}{(m+z)^s}\sum_{k=0}^{r-1}(-l)^k\binom{r-1}{k}m^{r-k-1}.$$

Thus, by the definition of $H_r(s,z)$, we have (2.36).
(2) First, we remark

$$\begin{aligned} K_r(s,z+1) &= \frac{1}{(r-1)!}\sum_{m\in\mathbb{Z}}\frac{(m)_{r-1}}{(m+z)^s} \\ &= \frac{1}{(r-1)!}\sum_{m\in\mathbb{Z}}\frac{(m+1)_r-(r-1)(m+1)_{r-1}}{(m+z)^s} \\ &= K_r(s,z) - K_{r-1}(s,z). \end{aligned}$$

Thus,

$$\begin{aligned} K_r(s, z+l) &= K_r(s, z+l-1) - K_{r-1}(s, z+l-1) \\ &= K_r(s,z) - \sum_{k=0}^{l-1} K_{r-1}(s, z+k). \end{aligned}$$

□

Proposition 2.12 (difference equations for s)**.** (1)

$$H_r(s-l, z) = \sum_{p=0}^{l} \binom{l}{p} z^{l-p} H_{r+p}(s, z). \tag{2.38}$$

(2)

$$K_r(s-l, z) = \frac{1}{(r-1)!} \sum_{k=1}^{r} \begin{bmatrix} r \\ k \end{bmatrix} \sum_{p=0}^{l} \binom{l}{p} z^{l-p} \sum_{q=1}^{k+p} (-1)^{k+p-q} (q-1)! \left\{ \begin{matrix} k+p \\ q \end{matrix} \right\} K_q(s, z). \tag{2.39}$$

Proof. (1) By the analytic continuation of $H_r(s, z)$, it is enough to show the assertion (2.38) when $\operatorname{Re} s > r + l$. In this case, it is easy to see that

$$\begin{aligned} H_r(s-l, z) &= \sum_{m \in \mathbb{Z}} \frac{m^{r-1}}{(m+z)^s} (m+z)^l \\ &= \sum_{m \in \mathbb{Z}} \frac{m^{r-1}}{(m+z)^s} \sum_{p=0}^{l} \binom{l}{p} z^{l-p} m^p \\ &= \sum_{p=0}^{l} \binom{l}{p} z^{l-p} \sum_{m \in \mathbb{Z}} \frac{m^{r+p-1}}{(m+z)^s} \\ &= \sum_{p=0}^{l} \binom{l}{p} z^{l-p} H_{r+p}(s, z). \end{aligned}$$

(2) From (2.13), (2.38) and (2.12), we have

$$\begin{aligned} K_r(s-l, z) &= \frac{1}{(r-1)!} \sum_{k=1}^{r} \begin{bmatrix} r \\ k \end{bmatrix} H_k(s-l, z) \\ &= \frac{1}{(r-1)!} \sum_{k=1}^{r} \begin{bmatrix} r \\ k \end{bmatrix} \sum_{p=0}^{l} \binom{l}{p} z^{l-p} H_{k+p}(s, z) \\ &= \frac{1}{(r-1)!} \sum_{k=1}^{r} \begin{bmatrix} r \\ k \end{bmatrix} \sum_{p=0}^{l} \binom{l}{p} z^{l-p} \sum_{q=1}^{k+p} (-1)^{k+p-q} (q-1)! \left\{ \begin{matrix} k+p \\ q \end{matrix} \right\} K_q(s, z). \end{aligned}$$

□

Proposition 2.13 (multiplication formulas)**.** (1)

$$H_r(s, Nz) = N^{-s} \sum_{k=0}^{N-1} \sum_{l=0}^{r-1} \binom{r-1}{l} k^l N^{r-l-1} H_{r-l}\left(s, z + \frac{k}{N}\right). \tag{2.40}$$

(2)

$$K_r(s, Nz) = N^{-s} \sum_{k_1,\ldots,k_r \geq 0}^{N-1} K_r\left(s, z + \frac{k_1 + \cdots + k_r}{N}\right). \tag{2.41}$$

Proof. (1) We decompose the sum of n by setting $n = Nj + k$.

$$\begin{aligned} H_r(s, Nz) &= \sum_{j\in\mathbb{Z}} \sum_{k=0}^{N-1} \frac{(Nj+k)^{r-1}}{(Nj+k+Nz)^s} \\ &= N^{-s+r-1} \sum_{k=0}^{N-1} \sum_{j\in\mathbb{Z}} \frac{1}{\left(j + z + \frac{k}{N}\right)^s} \sum_{l=0}^{r-1} \binom{r-1}{l} \left(\frac{k}{N}\right)^l j^{r-l-1} \\ &= N^{-s} \sum_{k=0}^{N-1} \sum_{l=0}^{r-1} \binom{r-1}{l} k^l N^{r-l-1} H_{r-l}\left(s, z + \frac{k}{N}\right). \end{aligned}$$

(2) First, we remark

$$\begin{aligned} \zeta_r(s, Nz) &= \sum_{j_1,\ldots,j_r\in\mathbb{Z}} \sum_{0\leq k_1,\ldots,k_r\leq N-1} \frac{1}{((Nj_1+k_1)+\ldots+(Nj_r+k_r)+Nz)^s} \\ &= N^{-s} \sum_{j_1,\ldots,j_r\in\mathbb{Z}} \sum_{0\leq k_1,\ldots,k_r\leq N-1} \frac{1}{\left(j_1+\ldots+j_r+z+\frac{k_1+\ldots+k_r}{N}\right)^s} \\ &= N^{-s} \sum_{0\leq k_1,\ldots,k_r\leq N-1} \zeta_r\left(s, z + \frac{k_1+\ldots+k_r}{N}\right) \end{aligned}$$

and

$$\begin{aligned} \zeta_r(s, r-Nz) &= \sum_{j_1,\ldots,j_r\in\mathbb{Z}} \sum_{1\leq k_1,\ldots,k_r\leq N} \frac{1}{((Nj_1+k_1)+\ldots+(Nj_r+k_r)-Nz)^s} \\ &= N^{-s} \sum_{j_1,\ldots,j_r\in\mathbb{Z}} \sum_{0\leq k_1,\ldots,k_r\leq N-1} \frac{1}{\left(j_1+\ldots+j_r+r-\left(z+\frac{k_1+\ldots+k_r}{N}\right)\right)^s} \\ &= N^{-s} \sum_{0\leq k_1,\ldots,k_r\leq N-1} \zeta_r\left(s, r - \left(z + \frac{k_1+\ldots+k_r}{N}\right)\right). \end{aligned}$$

Hence, by using (2.25),

$$\begin{aligned}
K_r(s,Nz) &= \zeta_r(s,Nz) + (-1)^{r-1}e^{-\pi is}\zeta_r(s,r-Nz) \\
&= N^{-s} \sum_{0\le k_1,\dots,k_r\le N-1} \\
&\quad \cdot \left\{ \zeta_r\left(s, z+\frac{k_1+\ldots+k_r}{N}\right) + (-1)^{r-1}e^{-\pi is}\zeta_r\left(s, r-\left(z+\frac{k_1+\ldots+k_r}{N}\right)\right)\right\} \\
&= N^{-s} \sum_{k_1,\dots,k_r\ge 0}^{N-1} K_r\left(s, z+\frac{k_1+\cdots+k_r}{N}\right).
\end{aligned}$$

□

Moreover, derivation formulas for $H_r(s,z)$ and $K_r(s,z)$ are obtained by easy calculations.

Proposition 2.14.

$$\frac{\partial}{\partial z}H_r(s,z) = -sH_r(s+1,z), \tag{2.42}$$

$$\frac{\partial}{\partial z}K_r(s,z) = -sK_r(s+1,z). \tag{2.43}$$

As a corollary of Proposition 2.14, we obtain the following important formulas, that is the logarithmic derivative of the generalized primitive and normalized multiple sine functions.

Corollary 2.15. (1)

$$\frac{\partial}{\partial z}\frac{\partial H_r}{\partial s}(0,z) = -H_r(1,z) = -(-z)^{r-1}(\pi i + \pi\cot\pi z), \tag{2.44}$$

$$\frac{\partial}{\partial z}\frac{\partial K_r}{\partial s}(0,z) = -K_r(1,z) = -\frac{(1-z)_{r-1}}{(r-1)!}(\pi i + \pi\cot\pi z). \tag{2.45}$$

(2) *For any $n>1$, we have*

$$\frac{\partial}{\partial z}\frac{\partial H_r}{\partial s}(1-n,z) = (n-1)\frac{\partial H_r}{\partial s}(2-n,z), \tag{2.46}$$

$$\frac{\partial}{\partial z}\frac{\partial K_r}{\partial s}(1-n,z) = (n-1)\frac{\partial K_r}{\partial s}(2-n,z). \tag{2.47}$$

Proof. Since the proofs for H and K are similar, we only give the proof for H.
(1) $H_r(s,z)$ is holomorphic around $s=0$ and $s=1$. Hence we have expressions

$$H_r(s,z) = \frac{\partial H_r}{\partial s}(0,z)s + O(s^2) = H_r(1,z) + O(s-1).$$

Thus,

$$\frac{\partial H_r}{\partial z}(s,z) = \frac{\partial}{\partial z}\frac{\partial H_r}{\partial s}(0,z)s + O(s^2).$$

On the other hands, from Proposition 2.14,

$$\frac{\partial H_r}{\partial z}(s,z) = -sH_r(s+1,z) = -H_r(1,z)s + O(s^2).$$

Therefore,

$$\frac{\partial}{\partial z}\frac{\partial H_r}{\partial s}(0,z) = -H_r(1,z) = -(-z)^{r-1}(\pi i + \pi\cot\pi z).$$

(2) For $n > 1$, $H_r(s,z)$ is holomorphic around $s = 1-n$ and $s = 2-n$. Hence we have expressions

$$H_r(s,z) = \frac{\partial H_r}{\partial s}(1-n,z)(s-1+n)+O((s-1+n)^2) = \frac{\partial H_r}{\partial s}(2-n,z)(s-2+n)+O((s-2+n)^2).$$

Thus,

$$\frac{\partial H_r}{\partial z}(s,z) = \frac{\partial}{\partial z}\frac{\partial H_r}{\partial s}(1-n,z)(s-1+n) + O((s-1+n)^2)$$

On the other hands, from Proposition 2.14,

$$\begin{aligned}\frac{\partial H_r}{\partial z}(s,z) &= -sH_r(s+1,z)\\ &= -(s-1+n+1-n)\left\{\frac{\partial H_r}{\partial s}(2-n,z)(s-1+n) + O((s-1+n)^2)\right\}\\ &= (n-1)\frac{\partial H_r}{\partial s}(2-n,z)(s-1+n) + O((s-1+n)^2).\end{aligned}$$

By comparing of $s-1+n$, we obtain the conclusion. □

3 A generalization of the multiple sine functions

Definition 3.1. We define the generalized primitive and normalized multiple sine functions as

$$\mathscr{S}_{r,n}(z) := \exp\left(-\frac{\partial H_r}{\partial s}(1-n,z)\right) \quad (\operatorname{Im} z > 0 \text{ or } 1 > z > -1), \tag{3.1}$$

$$\mathcal{S}_{r,n}(z) := \exp\left(-\frac{\partial K_r}{\partial s}(1-n,z)\right) \quad (\operatorname{Im} z > 0 \text{ or } r > z > 0). \tag{3.2}$$

If $n > 1$, $\mathcal{S}_{r,n}(z)$ is defined on $\operatorname{Im} z > 0$ or $r > z \geq 0$. Unless otherwise stated, we assume the above conditions of z.

From the definitions of $\mathscr{S}_{r,n}(z), \mathcal{S}_{r,n}(z)$ and the previous results, we obtain the following propositions immediately.

Proposition 3.2.

$$\mathscr{S}_{r,n}(z) = \prod_{k=1}^{r} \mathcal{S}_{k,n}(z)^{(-1)^{r-k}(k-1)!\left\{ {r \atop k} \right\}}, \tag{3.3}$$

$$\mathcal{S}_{r,n}(z) = \prod_{k=1}^{r} \mathscr{S}_{k,n}(z)^{\frac{1}{(r-1)!}\left[{r \atop k} \right]}. \tag{3.4}$$

Proposition 3.3. *Let $n \geq 1$.*

$$\mathscr{S}_{r,n}(z) = \exp\left(-\sum_{k=0}^{r-1} \frac{(n+k-1)!}{(2\pi i)^{n+k-1}} \binom{r-1}{k} (-z)^{r-1-k} \mathrm{Li}_{n+k}(e^{2\pi i z}) \right), \tag{3.5}$$

$$\mathcal{S}_{r,n}(z) = \exp\left(-\frac{1}{(r-1)!} \sum_{k=1}^{r} \begin{bmatrix} r \\ k \end{bmatrix} \sum_{l=0}^{k-1} \binom{k-1}{l} \frac{(n+l-1)!}{(2\pi i)^{n+l-1}} (-z)^{k-l-1} \mathrm{Li}_{n+l}(e^{2\pi i z}) \right). \tag{3.6}$$

Proposition 3.4. (1)

$$\mathscr{S}_{r,n}(0) = \exp\left(-\frac{(n+r-2)!}{(2\pi i)^{n+r-2}} \zeta(n+r-1) \right), \tag{3.7}$$

$$\mathcal{S}_{r,n+1}(0) = \prod_{k=1}^{r} \exp\left(-\frac{1}{(r-1)!} \begin{bmatrix} r \\ k \end{bmatrix} \frac{(n+k-1)!}{(2\pi i)^{n+k-1}} \zeta(n+k) \right). \tag{3.8}$$

(2)

$$\mathscr{S}_{r,n}\left(\frac{1}{2}\right) = \prod_{k=0}^{r-1} \exp\left(-\frac{(n+k-1)!}{(2\pi i)^{n+k-1}} \binom{r-1}{k} (-2)^{-r+1+k} (2^{1-n-k}-1) \zeta(n+k) \right), \tag{3.9}$$

$$\mathcal{S}_{r,n}\left(\frac{1}{2}\right) = \prod_{k=1}^{r} \prod_{l=0}^{k-1} \exp\left(-\frac{1}{(r-1)!} \begin{bmatrix} r \\ k \end{bmatrix} \binom{k-1}{l} \frac{(n+l-1)!}{(2\pi i)^{n+l-1}} (-2)^{-k+l+1} (2^{1-n-l}-1) \zeta(n+l) \right). \tag{3.10}$$

We remark $\lim_{n+k\to 1}(2^{1-n-k}-1)\zeta(n+k) = -\log 2$.

Proposition 3.5.

$$\mathscr{S}_{r,n}(z+l) = \prod_{k=0}^{r-1} \mathscr{S}_{r-k,n}(z)^{(-l)^k \binom{r-1}{k}}, \tag{3.11}$$

$$\mathcal{S}_{r,n}(z+l) = \mathcal{S}_{r,n}(z) \prod_{k=0}^{l-1} \mathcal{S}_{r-1,n}(z+k)^{-1}. \tag{3.12}$$

Proposition 3.6.

$$\mathscr{S}_{r,n}(z) = \prod_{p=0}^{n-1} \exp\left(\binom{n-1}{p} z^{n-p-1} \log \mathscr{S}_{r+p,1}(z) \right), \tag{3.13}$$

$$\mathcal{S}_{r,n}(z) = \prod_{k=1}^{r} \prod_{p=0}^{n-1} \prod_{q=1}^{k+p} \exp\left((-1)^{k+p-q} \frac{(q-1)!}{(r-1)!} \begin{bmatrix} r \\ k \end{bmatrix} \binom{n-1}{p} \left\{ \begin{matrix} k+p \\ q \end{matrix} \right\} z^{n-1-p} \log \mathcal{S}_{q,1}(z) \right). \tag{3.14}$$

Proposition 3.7.

$$\mathscr{S}_{r,n}(Nz) = \prod_{k=0}^{N-1} \prod_{l=0}^{r-1} \mathscr{S}_{r-l,n}\left(z + \frac{k}{N} \right)^{\binom{r-1}{l} k^l N^{n+r-l-2}}, \tag{3.15}$$

$$\mathcal{S}_{r,n}(Nz) = \prod_{k_1,\ldots,k_r \geq 0}^{N-1} \mathcal{S}_r\left(z + \frac{k_1 + \cdots + k_r}{N} \right)^{N^{n-1}}. \tag{3.16}$$

Proposition 3.8. (1)

$$\frac{\mathscr{S}'_{r,1}(z)}{\mathscr{S}_{r,1}(z)} = (-z)^{r-1}(\pi i + \pi \cot \pi z), \tag{3.17}$$

$$\frac{\mathcal{S}'_{r,1}(z)}{\mathcal{S}_{r,1}(z)} = \frac{(1-z)_{r-1}}{(r-1)!}(\pi i + \pi \cot \pi z). \tag{3.18}$$

(2) *For any* $n \in \mathbb{Z}_{\geq 1}$,

$$\frac{\mathscr{S}'_{r,n+1}(z)}{\mathscr{S}_{r,n+1}(z)} = n \log \mathscr{S}_{r,n}(z), \tag{3.19}$$

$$\frac{\mathcal{S}'_{r,n+1}(z)}{\mathcal{S}_{r,n+1}(z)} = n \log \mathcal{S}_{r,n}(z). \tag{3.20}$$

Consequently we have

$$\mathscr{S}_{r,n+1}(z) = \exp\left(-\frac{(n+r-1)!}{(2\pi i)^{n+r-1}} \zeta(n+r) \right) \exp\left(n \int_0^z \log \mathscr{S}_{r,n}(t)\, dt \right), \tag{3.21}$$

$$\mathcal{S}_{r,n+1}(z) = \exp\left(-\frac{1}{(r-1)!} \sum_{k=1}^{r} \begin{bmatrix} r \\ k \end{bmatrix} \frac{(n+k-1)!}{(2\pi i)^{n+k-1}} \zeta(n+k) \right) \exp\left(n \int_0^z \log \mathcal{S}_{r,n}(t)\, dt \right). \tag{3.22}$$

Proposition 3.9. *For any* $l \in \mathbb{Z}_{\geq 1}$,

$$n \int_0^l \log \mathscr{S}_{r,n}(z+t)\, dt = \sum_{k=1}^{r-1} (-l)^k \binom{r-1}{k} \log \mathscr{S}_{r-k,n+1}(z), \tag{3.23}$$

$$n \int_0^l \log \mathcal{S}_{r,n}(z+t)\, dt = -\sum_{k=0}^{l-1} \log \mathcal{S}_{r-1,n+1}(z+k). \tag{3.24}$$

Actually, Proposition 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 3.8 follow from Proposition 2.4, 2.9, Corollary 2.10, Proposition 2.11, 2.12, 2.13 and Corollary 2.15 respectively.

For (3.23),

$$\begin{aligned} n\int_0^l \log \mathscr{S}_{r,n}(z+t)\,dt &= \int_0^l \frac{d}{dt}\log \mathscr{S}_{r,n+1}(z+t)\,dt \\ &= \log \mathscr{S}_{r,n+1}(z+l) - \log \mathscr{S}_{r,n+1}(z) \\ &= \sum_{k=1}^{r-1}(-l)^k \binom{r-1}{k} \log \mathscr{S}_{r-k,n+1}(z). \end{aligned}$$

The first equality follows from (3.20) and the third one follows from (3.11). The proof of (3.24) can be similarly.

Furthermore, we also obtain the relations between our generalized multiple sine functions and known multiple sine function of [KK], [KW].

Proposition 3.10. (1)

$$\mathscr{S}_{r,1}(z) = \exp\left(-\frac{(r-1)!}{(2\pi i)^{r-1}}\zeta(r) + (-1)^{r-1}\pi i\frac{z^r}{r}\right)\mathscr{S}_r(z)^{(-1)^{r-1}}. \tag{3.25}$$

(2)

$$\mathcal{S}_{r,n}(z) = \exp\left((-1)^r\pi i\frac{(n-1)!}{(r+n-1)!}B_{r,r+n-1}(z)\right)S_{r,n}(z). \tag{3.26}$$

Here, $B_{r,k}(z)$ is the multiple Bernoulli polynomials defined by a generating function as

$$\frac{t^r e^{zt}}{(e^t-1)^r} = \sum_{k=0}^{\infty} B_{r,k}(z)\frac{t^k}{k!}.$$

Proof. (1) The logarithmic derivative of $\mathscr{S}_{r,1}(z)$ is

$$\frac{\mathscr{S}'_{r,1}(z)}{\mathscr{S}_{r,1}(z)} = (-z)^{r-1}(\pi i + \pi\cot\pi z) = (-1)^{r-1}\pi i z^{r-1} + (-1)^{r-1}\frac{\mathscr{S}'_r(z)}{\mathscr{S}_r(z)}.$$

The first equality follows from (2.44) and the second one from the integral expression of $\mathscr{S}_r(z)$

$$\mathscr{S}_r(z) = \exp\left(\int_0^z \pi t^{r-1}\cot(\pi t)\,dt\right).$$

Thus, there exists some constant C such that

$$\mathscr{S}_{r,1}(z) = C\exp\left((-1)^{r-1}\pi i\frac{z^r}{r}\right)\mathscr{S}_r(z)^{(-1)^{r-1}}.$$

In addition, by putting $z=0$,

$$C = \mathscr{S}_{r,1}(0).$$

(2) From the $\zeta_r(s,z)$ expression of $K_r(s,z)$ (2.25),

$$\begin{aligned}\mathcal{S}_{r,n}(z) &= \exp\left(-\frac{\partial K_r}{\partial s}(1-n,z)\right)\\ &= \exp\left((-1)^{r+n-1}\pi i\zeta_r(1-n,r-z) - \frac{\partial \zeta_r}{\partial s}(1-n,z) + (-1)^{r+n-1}\frac{\partial \zeta_r}{\partial s}(1-n,r-z)\right)\\ &= \exp\left((-1)^r\pi i\frac{(n-1)!}{(r+n-1)!}B_{r,r+n-1}(z)\right)S_{r,n}(z).\end{aligned}$$

The third equality follows from special values of $\zeta_r(1-n,X)$

$$\zeta_r(1-n,X) = (-1)^r\frac{(n-1)!}{(n+r-1)!}B_{r,r+n-1}(X)$$

and some formula of multiple Bernoulli polynomial

$$B_{r,k}(r-X) = (-1)^k B_{r,k}(X),$$

(see [B]). □

References

[AAR] G. E. Andrews, R. Askey and R. Roy: *Special Functions*, Encyclopedia of Mathematics and its Applications, 71. Cambridge University Press, 1999.

[B] E. W. Barnes: *On the theory of the multiple gamma function*, Trans. Cambridge Philos. Soc. **19** (1904), 374–425.

[KK] N. Kurokawa and S. Koyama: *Multiple sine functions*, Forum math. **15**-6 (2003), 839–876.

[KW] N. Kurokawa and M. Wakayama: *Period deformations and Raabe's formulas for generalized gamma and sine functions*, Kyushu J. Math. **62**-1 (2008), 171–187.

[O] K. Onodera: *Central values of generalized multiple sine functions*, Forum math. **21**-6 (2009), 1053–1065

[R] G. Rzadkowski: *Derivatives and Eulerian numbers*, Amer. Math. Monthly **115**-5 (2008), 458–460.

Institute of Mathematics for Industry, Kyushu University
744, Motooka, Nishi-ku, Fukuoka, 819-0395, JAPAN.
E-mail: g-shibukawa@math.kyushu-u.ac.jp